

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

## PATENT ABSTRACTS OF JAPAN

(11) Publication number : 10-306078  
 (43) Date of publication of application : 17.11.1998

(51) Int.Cl. C07D233/64  
 A61K 31/415  
 A61K 31/425  
 A61K 31/44  
 A61K 31/495  
 A61K 31/505  
 A61K 31/535  
 C07D213/30  
 C07D213/71  
 C07D213/74  
 C07D231/12  
 C07D233/61  
 C07D233/84  
 C07D239/42  
 C07D249/08  
 C07D277/36  
 C07D295/12  
 C07D307/64  
 C07D333/34  
 C07D401/12  
 C07D401/12  
 C07D401/12  
 C07D401/12  
 C07D403/12  
 C07D403/12

(21) Application number : 09-117976 (71) Applicant : MITSUBISHI CHEM CORP  
 (22) Date of filing : 08.05.1997 (72) Inventor : INOUE SHINYA





TARAO YOSHIHIRO  
 KOMATSU YOSHIYUKI  
 SUZUKI KAZUO  
 TAKAHASHI CHIZUKO

## (54) UREA DERIVATIVE

## (57) Abstract:

PROBLEM TO BE SOLVED: To obtain a new urea derivative useful as an active ingredient of a medicine used for prevention and/or treatment of diseases such as hyperlipemia or arteriosclerosis.

SOLUTION: This urea derivative is represented by formula I [R1 to R3 are each H, OH, an alkyl, an alkoxy, an aralkyloxy, etc.; R4 is H, an alkyl, a cycloalkyl, etc.; R5 to R7 are each H, an alkoxy, an alkyl, OH, etc.; Y is an alkyl, an aryl, etc.; (k) is 0-3; (1) is 2-4], its salt, hydrate or solvate, e.g. N-[4-(4-methyl-1-piperazinyl)phenyl]methyl-N'-[2-[3-(4-phenyl-1-piperazinyl) propoxy]-6-methylphenyl]urea. The compound represented by formula I is obtained by converting an aniline derivative represented by formula II into a reactional intermediate represented by formula III



(Z is a halogen, an aryloxy, etc.), then reacting the resultant intermediate with an amine derivative represented by formula IV in which (k) is 1-3 or an aniline derivative represented by formula IV in which (k) is 0.

---

**LEGAL STATUS**

[Date of request for examination]

[Date of sending the examiner's decision of rejection]

[Kind of final disposal of application other than the examiner's decision of rejection or application converted registration]

[Date of final disposal for application]

[Patent number]

[Date of registration]

[Number of appeal against examiner's decision of rejection]

[Date of requesting appeal against examiner's decision of rejection]

[Date of extinction of right]

Copyright (C); 1998,2000 Japan Patent Office